

Lymphoma scanning

		Interim scan	End of chemo	Post RT/ RM
Burkitt	No routine interim scanning in patients with clinical response	If response not assessable: -CT post 3x DA-EPOCH or 2x R-CODOX/M-IVAC -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo* (CT if prior CMR)	
DLBCL advanced	Routine iPET-2 Chemo to be continued without awaiting result	All patients: PET D15-20 of C2 R-CHOP/pola-R-CHP PET after C3 if PET-2 not feasible	PET scan 6 weeks after last dose of chemo**	If RT: PET 3 months post RT (CT if prior CMR)
DLBCL For CMT	Routine iPET-2 Chemo to be continued without awaiting result	All patients: PET D15-20 of C2 R-CHOP/pola-R-CHP	RT to <i>start</i> within 3-5 weeks of last chemo No repeat scan if iPET-2 done	PET 3 months post RT (CT if prior CMR)
PMBL R-CHOP	Routine iPET-2 Chemo to be continued without awaiting result	All patients: PET D10-13 of C2 R-CHOP-14 PET after C3 if PET-2 not feasible	No pre-RT scan if PET-2= CMR Others: PET 3 weeks after last dose (RT to <i>start</i> 3-6w post)	PET 3 months post RT (CT if prior CMR)
PMBL DA-EPOCH	Routine iPET-2 Chemo to be continued without awaiting result	All patients: PET D15-20 of C2 DA-EPOCH-R PET after C3 if PET-2 not feasible	PET scan 4 weeks after last dose of chemo (CT if prior CMR)	If RT (EOT PMR): PET 3 months post RT
FL	No routine interim scanning in patients with clinical response	If response not assessable: -CT post 4x R-chemo -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo (no EOT scan if prior CMR)	PET scan at 12 months of RM (CT if prior CMR)
MZL	No routine interim scanning in patients on R-chemo Routine CT after 4x R mono	R-chemo: if response not assessable: -CT post 4x R-chemo R mono: CT 4 weeks after 4 x weekly	CT scan 4 weeks after last dose of chemo/R mono (no EOT scan if prior CR)	CT at 12 months of RM
MCL	No routine interim scanning in patients with clinical response unless auto-SCT planned	If response not assessable/planned SCT: -CT post 3x R-chemo -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo^ (CT if prior CMR)	PET scan at 12 months of RM (CT if prior CMR)

*disregard HD-MTX or intrathecal

**3-4 weeks if HD MTX at EOT, or pre-planned consolidation RT not driven by PET response

^3 weeks if SCT is planned